



Scattering Amplitudes in Maximally Supersymmetric Gauge Theory and a New Duality



Lance Dixon (SLAC)

LD, Ö. Gürdoğan, A. McLeod, M. Wilhelm, 2112.06243


HEP Seminar
Northwestern University
Valentines Day, 2022



CMS Experiment at LHC, CERN
Data recorded: Mon Oct 25 05:47:22 2010 CDT
Run/Event: 148844 / 19210956
Lumi section: 520
Orbit/Crossing: 136152948 / 1594

LHC is a QCD Machine


- Copious production of quarks and gluons which materialize as collimated jets of hadrons



- Confrontation between experiment and theory at high precision requires taking into account **higher order corrections** in the **strong coupling α_s**

Example: Total cross section for producing Higgs boson at LHC via gluon fusion

Leading Order (LO)



- Higgs production at LHC is dominantly via gluon fusion, mediated by a top quark loop.
- Since $2m_{top} = 350 \text{ GeV}$
 $\gg m_{Higgs} = 125 \text{ GeV}$,
 we can integrate out the top quark to get a leading operator $H G_{\mu\nu}^a G^{\mu\nu a}$


Perturbative Short-Distance Cross Section

$$\hat{\sigma}(\alpha_s, \mu_F, \mu_R) = [\alpha_s(\mu_R)]^{n_\alpha} \left[\underbrace{\hat{\sigma}^{(0)}}_{\text{LO}} + \frac{\alpha_s}{2\pi} \underbrace{\hat{\sigma}^{(1)}}_{\text{NLO}}(\mu_F, \mu_R) + \left(\frac{\alpha_s}{2\pi}\right)^2 \underbrace{\hat{\sigma}^{(2)}}_{\text{NNLO}}(\mu_F, \mu_R) + \dots \right]$$


Leading-order (LO) predictions **qualitative**: **poor convergence** of expansion in $\alpha_s(\mu)$
 Uncertainty bands from varying $\mu_R = \mu_F = \mu$

Example: Higgs gluon fusion cross section at LHC vs. CM energy \sqrt{s}


LO \rightarrow NNNLO \rightarrow factor of 2.7 increase!



NLO QCD topologies




virtual $gg \rightarrow H$



real, $gg \rightarrow Hg$

NNNLO QCD topologies



Multi-loop complexities

- Multi-loop multiscale integrals typically very difficult to evaluate
- All 1 loop integrals reducible to scalar box integrals + simpler

→ combinations of
+ simpler

$$\text{Li}_2(x) = - \int_0^x \frac{dt}{t} \ln(1-t)$$

Brown-Feynman (1952), Melrose (1965), 't Hooft-Veltman (1974), Passarino-Veltman (1979), van Neerven-Vermaseren (1984), Bern, LD, Kosower (1992)

- At L loops, special functions with up to $2L$ integrations
Hannesdottir, McLeod, Schwartz, Vergu, 2109.09744
- Weight $2L$ iterated integrals, generalized polylogarithms, or worse

Planar N=4 SYM, toy model for QCD amplitudes

- QCD's maximally supersymmetric cousin, N=4 super-Yang-Mills theory (SYM), gauge group $SU(N_c)$, in large N_c (planar) limit
- Structure very rigid:
Amplitudes = $\sum_i \text{rational}_i \times \text{transcendental}_i$
- For planar N=4 SYM, we understand rational structure quite well, focus on the transcendental functions.
- Furthermore, at least three dualities hold:
 1. AdS/CFT
 2. Amplitudes dual to Wilson loops
 3. New “antipodal” duality between amplitudes and form factors

Finite radius of convergence

- Planar N=4 SYM has **no renormalons** ($\beta(g) = 0$) and **no instantons** ($e^{-1/g_{\text{YM}}^2} = e^{-N_c/\lambda}$)
- Its perturbative expansion can have a **finite radius of convergence**, unlike QCD, QED, whose perturbative series are **asymptotic**.
- For cusp anomalous dimension, using coupling

$$g^2 \equiv \frac{N_c g_{\text{YM}}^2}{16\pi^2} = \frac{\lambda}{16\pi^2}, \quad \text{the radius is } \frac{1}{16}$$

Beisert, Eden, Staudacher (BES), 0610251

- Ratio of successive loop orders $\frac{\Gamma_{\text{cusp}}^{(L)}}{\Gamma_{\text{cusp}}^{(L-1)}} \rightarrow -16$
- Evidence for **same radius of convergence in high-loop-order behavior of amplitudes and form factors**, in suitable kinematic regions.

N=4 SYM particle content

Brink, Schwarz, Scherk; Gliozzi, Scherk, Olive (1977)

massless spin 1 gluon 
 4 massless spin 1/2 gluinos 
 6 massless spin 0 scalars 

$$G = SU(N_c)$$

SUSY
 $Q_a, a=1,2,3,4$
 shifts helicity
 by $1/2 \leftrightarrow$


$\mathcal{N} = 4$	1	\leftrightarrow	4	\leftrightarrow	6	\leftrightarrow	4	\leftrightarrow	1
	g^-		$\lambda_{\bar{i}}^-$		$\bar{\phi}_{\bar{i}\bar{j}}, \phi_{ij}$		λ_i^+		g^+
helicity	-1		$-\frac{1}{2}$		0		$\frac{1}{2}$		1

all in adjoint representation of G





QCD vs. N=4 SYM

- QCD has **gluons** and **quarks** in fundamental rep. of $SU(N_c)$
- Replace **quarks** with 4 copies of fermions in adjoint rep. (**gluinos**) and add 6 real adjoint **scalars**
- Feynman vertices:

QCD





N=4 SYM



QCD vs. N=4 SYM at tree level

At tree-level essentially identical

Consider a tree amplitude for n gluons.
Fermions and **scalars** cannot appear
 because they are produced in **pairs**



Hence the amplitude is the same in QCD and N=4 SYM.
 The QCD tree amplitude “secretly” obeys **all identities of N=4 supersymmetry**:

$$\begin{aligned}
 & \text{Diagram 1} = \text{Diagram 2} = 0 \quad \frac{1}{\langle ij \rangle^4} \times \text{Diagram 3} \quad \text{independent of } i, j
 \end{aligned}$$

The diagrams show a central brown oval with four green wavy lines extending from it. In the first diagram, all four lines have a '+' sign. In the second, the top two have '+' and the bottom two have '-'. In the third, the top-left and bottom-right have '+', while the top-right and bottom-left have '-'.


No longer true at quantum (loop) level

N=4 SYM very special

- At one loop, cancellation of loop momenta in numerator
 \rightarrow only scalar box integrals



Bern, LD, Dunbar, Kosower, hep-ph/9403226

- Weight 2 functions – dilogs. E.g., $gg \rightarrow Hg$ @ 1 loop \supset



$$= \text{Li}_2\left(1 - \frac{s_{123}}{s_{12}}\right) + \text{Li}_2\left(1 - \frac{s_{123}}{s_{23}}\right) + \frac{1}{2} \ln^2\left(\frac{s_{12}}{s_{23}}\right) + \dots$$

- QCD results also contain single log's and rational parts from (tensor) triangle + bubble integrals

$$= \frac{1}{\epsilon} - \ln(s_{123})$$

Higher loops

- Much evidence that N=4 SYM amplitudes have “uniform **weight** (transcendentality)” $2L$ at loop order L
- **Weight** \sim number of integrations, e.g.

$$\ln(s) = \int_1^s \frac{dt}{t} = \int_1^s d\ln t \quad 1$$


$$\text{Li}_2(x) = - \int_0^x \frac{dt}{t} \ln(1-t) = \int_0^x d\ln t \cdot [-\ln(1-t)] \quad 2$$

$$\text{Li}_n(x) = \int_0^x \frac{dt}{t} \text{Li}_{n-1}(t) \quad n$$

AdS/CFT

Maldacena (1997)

Conformal field theory (like N=4 SYM) is dual to a theory of gravity in anti-de Sitter space (like strings in $AdS_5 \times S^5$)



SO(4,2) isometry
of 5 dimensional
space-time

\leftrightarrow 4d conformal symmetry

A weak-strong duality

T-duality symmetry of string theory

Alday, Maldacena, 0705.0303

- Exchanges string world-sheet variables σ, τ

- $X^\mu(\tau, \sigma) = x^\mu + k^\mu \tau + \text{oscillators}$

$\rightarrow X^\mu(\tau, \sigma) = x^\mu + k^\mu \sigma + \text{oscillators}$



- **Strong coupling** limit of planar N=4 SYM

is **semi-classical** limit of string theory:


world-sheet stretches tight around

minimal area surface in AdS.

- Boundary determined by **momenta** of external states: **light-like polygon with null edges = momenta k^μ**



Amplitudes = Wilson loops



- Polygon vertices x_i are not positions but **dual momenta**,
$$x_i - x_{i+1} = k_i$$
- Transform like positions under **dual conformal symmetry**

Alday, Maldacena, 0705.0303
Drummond, Korchemsky, Sokatchev, 0707.0243
Brandhuber, Heslop, Travaglini, 0707.1153
Drummond, Henn, Korchemsky, Sokatchev,
0709.2368, 0712.1223, 0803.1466;
Bern, LD, Kosower, Roiban, Spradlin,
Vergu, Volovich, 0803.1465

Duality verified to hold
at weak coupling too

weak-weak duality,
holds order-by-order

Dual conformal invariance

- Wilson n -gon invariant under inversion: $x_i^\mu \rightarrow \frac{x_i^\mu}{x_i^2}$, $x_{ij}^2 \rightarrow \frac{x_{ij}^2}{x_i^2 x_j^2}$
 $x_{ij}^2 = (k_i + k_{i+1} + \dots + k_{j-1})^2 \equiv s_{i,i+1,\dots,j-1}$

- Fixed, up to functions of invariant cross ratios:

$$u_{ijkl} \equiv \frac{x_{ij}^2 x_{kl}^2}{x_{ik}^2 x_{jl}^2}$$


- $x_{i,i+1}^2 = k_i^2 = 0 \rightarrow$ no such variables for $n = 4, 5$

$n = 6 \rightarrow$ precisely 3 ratios:

$n = 7 \rightarrow$ 6 ratios.

In general, $3n-15$ ratios.

$$\left. \begin{aligned} u &= \frac{x_{13}^2 x_{46}^2}{x_{14}^2 x_{36}^2} = \frac{s_{12} s_{45}}{s_{123} s_{345}} \\ v &= \frac{s_{23} s_{56}}{s_{234} s_{123}} \\ w &= \frac{s_{34} s_{61}}{s_{345} s_{234}} \end{aligned} \right\}$$



Hexagon function bootstrap

Loops

3

LD, Drummond, Henn, 1108.4461, 1111.1704;


4,5

Caron-Huot, LD, Drummond, Duhr, von Hippel, McLeod, Pennington, 1308.2276, 1402.3300, 1408.1505, 1509.08127; 1609.00669;

6,7


Caron-Huot, LD, Dulat, von Hippel, McLeod, Papathanasiou, 1903.10890, 1906.07116; LD, Dulat, 22mm.nnnnn (NMHV 7 loop)

- Use analytical properties of perturbative (six point) amplitudes in planar N=4 SYM to determine them directly, **without ever peeking inside the loops**
- Step toward doing this **nonperturbatively (no loops to peek inside)** for general kinematics



6-gluon kinematics:
many different regions


Multi-particle
factorization $u, w \rightarrow \infty$



Example: MHV finite remainder $R_6^{(L)}$ on (u,u,u)

Alday, Gaiotto,
Maldacena,
0911.4708

$$R_6^{(L)}(u,u,u)/R_6^{(L)}(1,1,1)$$



- **Amazing proportionality** of each perturbative coefficient at small u , and also with the strong coupling result

Origin at weak coupling

- Remarkably, MHV remainder R_6 and closely-related quantity $\ln \mathcal{E}$ are **quadratic in logarithms** through 7 loops CDDvHMP, 1903.10890
- Previously observed through 2 loops, and at strong coupling, on the diagonal (u, u, u) AGM, 0911.4708

$$\ln \mathcal{E}(u_i) \approx -\frac{\Gamma_{\text{oct}}}{24} \ln^2(u_1 u_2 u_3) - \frac{\Gamma_{\text{hex}}}{24} \sum_{i=1}^3 \ln^2 \frac{u_i}{u_{i+1}} + C_0$$


	$L = 1$	$L = 2$	$L = 3$	$L = 4$	$L = 5$
Γ_{oct}	4	$-16\zeta_2$	$256\zeta_4$	$-3264\zeta_6$	$\frac{126976}{3}\zeta_8$
Γ_{cusp}	4	$-8\zeta_2$	$88\zeta_4$	$-876\zeta_6 - 32\zeta_3^2$	$\frac{28384}{3}\zeta_8 + 128\zeta_2\zeta_3^2 + 640\zeta_3\zeta_5$
Γ_{hex}	4	$-4\zeta_2$	$34\zeta_4$	$-\frac{603}{2}\zeta_6 - 24\zeta_3^2$	$\frac{18287}{6}\zeta_8 + 48\zeta_2\zeta_3^2 + 480\zeta_3\zeta_5$
C_0	$-3\zeta_2$	$\frac{77}{4}\zeta_4$	$-\frac{4463}{24}\zeta_6 + 2\zeta_3^2$	$\frac{67645}{32}\zeta_8 + 6\zeta_2\zeta_3^2 - 40\zeta_3\zeta_5$	$-\frac{4184281}{160}\zeta_{10} - 65\zeta_4\zeta_3^2 - 120\zeta_2\zeta_3\zeta_5 + 228\zeta_5^2 + 420\zeta_3\zeta_7$

- Coefficients involve same **BES kernel** as for **cusp**, but “tilted” by angle α ,
 $\Gamma_{\text{cusp}} = \Gamma_{\alpha=\pi/4}$ $\Gamma_{\text{oct}} = \Gamma_{\alpha=0}$ $\Gamma_{\text{hex}} = \Gamma_{\alpha=\pi/3}$

B. Basso, LD, G. Papathanasiou, 2001.05460

Solving for Planar N=4 SYM Amplitudes


Images: A. Sever, N. Arkani-Hamed




“Higgs” amplitudes and N=4 SYM form factors

LD, A. McLeod, M. Wilhelm, 2012.12286
+ Ö. Gürdoğan, to appear

3,4,5 loops
6,7,8 loops



- At leading order in $1/m_{top}$, Higgs boson couples to gluons via the operator $H G_{\mu\nu}^a G^{\mu\nu a}$



Form factors (cont.)

- Higgs is scalar, color singlet. QCD amplitudes with gluons are matrix elements of $G_{\mu\nu}^a G^{\mu\nu a}$ with on-shell gluons: “form factors”
- In N=4, operator is part of (BPS-protected) stress tensor supermultiplet, also includes for example $\phi_1^\dagger \phi_1 - \phi_2^\dagger \phi_2$ ($\in \mathbf{20}$ of $SU(4)_R$)
- Hgg “Sudakov” form factor is “too simple”; no kinematic dependence beyond overall $(-s_{12})^{-L\epsilon}$
- $Hggg$ is “just right”, depends on 2 dimensionless ratios


Hggg kinematics is two-dimensional

$$k_1 + k_2 + k_3 = -k_H$$

$$s_{123} = s_{12} + s_{23} + s_{31} = m_H^2$$

$$s_{ij} = (k_i + k_j)^2 \quad k_i^2 = 0$$

$$u = \frac{s_{12}}{s_{123}} \quad v = \frac{s_{23}}{s_{123}} \quad w = \frac{s_{31}}{s_{123}}$$



$$u + v + w = 1$$

I = decay / Euclidean

IIa,b,c = scattering / spacelike operator

IIIa,b,c = scattering / timelike operator

not cross ratios!

N=4 amplitude is S_3 invariant


$D_3 \equiv S_3$ dihedral symmetry generated by:

a. cycle: $i \rightarrow i + 1 \pmod{3}$, or

$$u \rightarrow v \rightarrow w \rightarrow u$$

b. flip: $u \leftrightarrow v$

One loop integrals/amplitudes



$$= \text{Li}_2\left(1 - \frac{s_{123}}{s_{12}}\right) + \text{Li}_2\left(1 - \frac{s_{123}}{s_{23}}\right) + \frac{1}{2} \ln^2\left(\frac{s_{12}}{s_{23}}\right) + \dots$$

$$= \text{Li}_2\left(1 - \frac{1}{u}\right) + \text{Li}_2\left(1 - \frac{1}{v}\right) + \frac{1}{2} \ln^2\left(\frac{u}{v}\right) + \dots$$

A two-loop story

- Gehrman et al. computed $Hggg$ in QCD at 2 loops
Gehrman, Jaquier, Glover, Koukoutsakis, 1112.3554
- Soon after, Brandhuber et al. computed stress tensor 3-point form factor \mathcal{F}_3 in N=4 SYM,
Brandhuber, Travaglini, Yang, 1201.4170
saw that “maximally transcendental part” of QCD result (both (+++) and (-++)) was **same as N=4 result!!**
- This “principle of maximal transcendentalty”
Kotikov, Lipatov, Velizhanin, hep-ph/0301021, hep-ph/0611204
was known to work for DGLAP and BFKL anomalous dimensions, but **not** for generic scattering amplitudes, so this one is **very special**

2d HPLs

Gehrmann, Remiddi, hep-ph/0008287

Space graded by weight n . Every function F obeys:

$$\frac{\partial F(u, v)}{\partial u} = \frac{F^u}{u} - \frac{F^w}{w} - \frac{F^{1-u}}{1-u} + \frac{F^{1-w}}{1-w}$$
$$\frac{\partial F(u, v)}{\partial v} = \frac{F^v}{v} - \frac{F^w}{w} - \frac{F^{1-v}}{1-v} + \frac{F^{1-w}}{1-w}$$

$$w = 1 - u - v$$

where $F^u, F^v, F^w, F^{1-u}, F^{1-v}, F^{1-w}$ are weight $n-1$ 2d HPLs.

To bootstrap $Hggg$ amplitude beyond 2 loops, find **as small a subspace of 2d HPLs as possible**, construct it to high weight.

Generalized polylogarithms

Chen, Goncharov, Brown,...

- Can be defined as **iterated integrals**, e.g.

$$G(a_1, a_2, \dots, a_n, x) = \int_0^x \frac{dt}{t - a_1} G(a_2, \dots, a_n, t)$$

- Or define differentially:

$$dF = \sum_{s_k \in \mathcal{S}} F^{s_k} d \ln s_k$$

- There is a Hopf algebra that “co-acts” on the space of polylogarithms, $\Delta: F \rightarrow F \otimes F$
- The **derivative** dF is one piece of Δ : $\Delta_{n-1,1} F = \sum_{s_k \in \mathcal{S}} F^{s_k} \otimes \ln s_k$
- so we refer to F^{s_k} as a $\{n-1,1\}$ coproduct of F
- s_k are letters in the symbol alphabet \mathcal{S}

Generalized polylogarithms (cont.)

- The $\{n-1,1\}$ coaction can be applied iteratively.
- Define the $\{n-2,1,1\}$ **double** coproducts, F^{S_k, S_j} , via the derivatives of the $\{n-1,1\}$ **single** coproducts F^{S_j} :

$$dF^{S_j} \equiv \sum_{S_k \in \mathcal{S}} F^{S_k, S_j} d \ln s_k$$

- And so on for the $\{n-m, 1, \dots, 1\}$ m^{th} coproducts of F .
- The **maximal iteration**, n times for a weight n function, is the **symbol**,

$$\mathcal{S}[F] = \sum_{S_{i_1}, \dots, S_{i_n} \in \mathcal{S}} F^{S_{i_1}, \dots, S_{i_n}} d \ln s_{i_1} \dots d \ln s_{i_n} \equiv \sum_{S_{i_1}, \dots, S_{i_n} \in \mathcal{S}} F^{S_{i_1}, \dots, S_{i_n}} s_{i_1} \otimes \dots \otimes s_{i_n}$$

where now $F^{S_{i_1}, \dots, S_{i_n}}$ are just rational numbers

Goncharov, Spradlin, Vergu, Volovich, 1006.5703

Example: Classical polylogarithms

$$\text{Li}_1(x) = -\ln(1-x) = \sum_{k=1}^{\infty} \frac{x^k}{k}$$

$$\text{Li}_n(x) = \int_0^x \frac{dt}{t} \text{Li}_{n-1}(t) = \sum_{k=1}^{\infty} \frac{x^k}{k^n}$$

- Regular at $x = 0$, branch cut starts at $x = 1$.
- Iterated differentiation gives the symbol:

$$\begin{aligned} \mathcal{S}[\text{Li}_n(x)] &= \mathcal{S}[\text{Li}_{n-1}(x)] \otimes x \\ &= \dots = -(1-x) \otimes x \otimes \dots \otimes x \end{aligned}$$

- **Branch cut** discontinuities displayed in **first** entry of symbol, e.g. clip off leading $(1-x)$ to compute discontinuity at $x = 1$.
- **Derivatives** computed from symbol by clipping **last** entry, multiplying by that $d \ln(\dots)$.

Example: Harmonic Polylogarithms in one variable (HPL{0,1})

Remiddi, Vermaseren, hep-ph/9905237

- Generalize the classical polylogs
- Define HPLs by iterated integration:

$$H_{0,\vec{w}}(x) = \int_0^x \frac{dt}{t} H_{\vec{w}}(t), \quad H_{1,\vec{w}}(x) = \int_0^x \frac{dt}{1-t} H_{\vec{w}}(t)$$

- Or by derivatives:

$$dH_{0,\vec{w}}(x) = H_{\vec{w}}(x) d \ln x \quad dH_{1,\vec{w}}(x) = -H_{\vec{w}}(x) d \ln(1-x)$$

- Symbol alphabet: $\mathcal{S} = \{x, 1-x\}$
- Weight n = length of binary string \vec{w}
- Number of functions at weight $n = 2L$ is number of binary strings: 2^{2L}
- **Branch cuts** dictated by **first** integration/entry in symbol
- **Derivatives** dictated by **last** integration/entry in symbol

Symbol alphabet \mathcal{S} for H_{ggg}

Gehrmann, Remiddi, hep-ph/0008287

- Comparing

$$\frac{\partial F(u, v)}{\partial u} = \frac{F^u}{u} - \frac{F^w}{w} - \frac{F^{1-u}}{1-u} + \frac{F^{1-w}}{1-w}$$

$$\frac{\partial F(u, v)}{\partial v} = \frac{F^v}{v} - \frac{F^w}{w} - \frac{F^{1-v}}{1-v} + \frac{F^{1-w}}{1-w}$$

$$w = 1 - u - v$$

with

$$dF = \sum_{s_k \in \mathcal{S}} F^{s_k} d \ln s_k$$

alphabet is $\mathcal{S} = \{u, v, w, 1 - u, 1 - v, 1 - w\}$

Heuristic view of function space

weight

...


4

3

2

1

0



$$\frac{\partial F(u, v)}{\partial u} = \frac{F^u}{u} - \frac{F^w}{w} - \frac{F^{1-u}}{1-u} + \frac{F^{1-w}}{1-w}$$

Symbol alphabets for n -gluon amplitudes

parity-odd letters, algebraic in $\hat{u}, \hat{v}, \hat{w}$

$n = 6$ has 9 letters: $\mathcal{S}_6 = \{\hat{u}, \hat{v}, \hat{w}, 1 - \hat{u}, 1 - \hat{v}, 1 - \hat{w}, \hat{y}_u, \hat{y}_v, \hat{y}_w\}$

Goncharov, Spradlin, Vergu, Volovich, 1006.5703;
LD, Drummond, Henn, 1108.4461; Caron-Huot,
LD, von Hippel, McLeod, 1609.00669



$n = 7$ has 42 letters

Golden, Goncharov, Paulos, Spradlin, Volovich, Vergu, 1305.1617,
1401.6446, 1411.3289; Drummond, Papathanasiou, Spradlin 1412.3763

$n = 8$ has at least 222 letters, could even be infinite as $L \rightarrow \infty$

Arkani-Hamed, Lam, Spradlin, 1912.08222;
Drummond, Foster, Gürdoğan, Kalousios, 1912.08217, 2002.04624;
Henke, Papathanasiou 1912.08254, 2106.01392;
Z. Li, C. Zhang, 2110.00350

Beyond $n = 8$



More bootstrapping details

- Four lectures at “The Amplitude Games”, Mainz Institute for Theoretical Physics, July, 2021
- <https://indico.mitp.uni-mainz.de/event/204/sessions/875/>

3-gluon form factor: better alphabet

- Motivated by 6 gluon case, switch to equivalent alphabet

$$\mathcal{S}' = \left\{ a = \frac{u}{vw}, b = \frac{v}{wu}, c = \frac{w}{uv}, d = \frac{1-u}{u}, e = \frac{1-v}{v}, f = \frac{1-w}{w} \right\}$$

- Symbols of (suitably normalized) form factor $F_3^{(L)}$ simplify remarkably at 1 and 2 loops, just 1 and 2 terms, plus D_3 dihedral images(!!!):

$$S \left[F_3^{(1)} \right] = (-1) b \otimes d + \text{dihedral}$$

$$S \left[F_3^{(2)} \right] = 4 b \otimes d \otimes d \otimes d + 2 b \otimes b \otimes b \otimes d + \text{dihedral}$$

dihedral cycle: $a \rightarrow b \rightarrow c \rightarrow a, \quad d \rightarrow e \rightarrow f \rightarrow d$

dihedral flip: $a \leftrightarrow b, \quad d \leftrightarrow e$

Simplest analytic form is for $v \rightarrow \infty$

→ Harmonic polylogarithms $H_{\vec{w}} \equiv H_{\vec{w}}(1 - \frac{1}{u})$

$$F_3^{(1)}(v \rightarrow \infty) = 2H_{0,1} + 6\zeta_2$$

$$F_3^{(2)}(v \rightarrow \infty) = -8H_{0,0,0,1} - 4H_{0,1,1,1} + 12\zeta_2 H_{0,1} + 13\zeta_4$$

$$\begin{aligned} F_3^{(3)}(v \rightarrow \infty) = & 96H_{0,0,0,0,0,1} + 16H_{0,0,0,1,0,1} + 16H_{0,0,0,1,1,1} + 16H_{0,0,1,0,0,1} + 8H_{0,0,1,0,1,1} \\ & + 8H_{0,0,1,1,0,1} + 16H_{0,1,0,0,0,1} + 8H_{0,1,0,0,1,1} + 12H_{0,1,0,1,0,1} + 4H_{0,1,0,1,1,1} \\ & + 8H_{0,1,1,0,0,1} + 4H_{0,1,1,0,1,1} + 4H_{0,1,1,1,0,1} + 24H_{0,1,1,1,1,1} \\ & - \zeta_2(32H_{0,0,0,1} + 8H_{0,0,1,1} + 4H_{0,1,0,1} + 52H_{0,1,1,1}) \\ & - \zeta_3(8H_{0,0,1} - 4H_{0,1,1}) - 53\zeta_4 H_{0,1} - \frac{167}{4}\zeta_6 + 2(\zeta_3)^2 \end{aligned}$$

8 loop result has $\sim 2^{2 \times 8 - 2} = 16,384$ terms

6-gluon amplitude is simplest for $(\hat{u}, \hat{v}, \hat{w}) = (1, \hat{v}, \hat{v})$

- Let $H_{\vec{w}} \equiv H_{\vec{w}}(1 - \frac{1}{\hat{v}})$

$$A_6^{(1)}(1, \hat{v}, \hat{v}) = 2H_{0,1}$$

$$A_6^{(2)}(1, \hat{v}, \hat{v}) = -8H_{0,1,1,1} - 4H_{0,0,0,1} - 4\zeta_2 H_{0,1} - 9\zeta_4$$

$$\begin{aligned} A_6^{(3)}(1, \hat{v}, \hat{v}) = & 96H_{0,1,1,1,1,1} + 16H_{0,1,0,1,1,1} + 16H_{0,0,0,1,1,1} + 16H_{0,1,1,0,1,1} + 8H_{0,0,1,0,1,1} \\ & + 8H_{0,1,0,0,1,1} + 16H_{0,1,1,1,0,1} + 8H_{0,0,1,1,0,1} + 12H_{0,1,0,1,0,1} + 4H_{0,0,0,1,0,1} \\ & + 8H_{0,1,1,0,0,1} + 4H_{0,0,1,0,0,1} + 4H_{0,1,0,0,0,1} + 24H_{0,0,0,0,0,1} \\ & + \zeta_2(8H_{0,0,0,1} + 8H_{0,1,0,1} + 48H_{0,1,1,1}) \\ & + 42\zeta_4 H_{0,1} + 121\zeta_6 \end{aligned}$$

Exact map at symbol level, with $\frac{1}{\hat{v}} = 1 - \frac{1}{u}$, $0 \leftrightarrow 1$,

if you also **reverse order** of symbol entries / HPL indices!!!

Works to **7 loops**, where $\sim 2^{2 \times 7 - 2} = 4,096$ terms agree

Antipodal duality in 2d

weak-weak duality

LD, Ö. Gürdoğan, A. McLeod,
M. Wilhelm, 2112.06243

$$F_3^{(L)}(u, v, w) = S \left(A_6^{(L)}(\hat{u}, \hat{v}, \hat{w}) \right)$$

Antipode map S , at symbol level, **reverses order of all letters**:

$$S(x_1 \otimes x_2 \otimes \cdots \otimes x_m) = (-1)^m x_m \otimes \cdots \otimes x_2 \otimes x_1$$

Kinematic map is

$$\hat{u} = \frac{vw}{(1-v)(1-w)}, \quad \hat{v} = \frac{wu}{(1-w)(1-u)}, \quad \hat{w} = \frac{uv}{(1-u)(1-v)}$$

Maps $u + v + w = 1$ to parity-preserving surface

$$\Delta \equiv (1 - \hat{u} - \hat{v} - \hat{w})^2 - 4\hat{u}\hat{v}\hat{w} = 0$$

also corresponds to “twisted forward scattering”:

$$\hat{k}_{i+n}^\mu = -\hat{k}_i^\mu, \quad i = 1, 2, \dots, n \quad (n = 3 \text{ here})$$

6-gluon alphabet and symbol map

- $\mathcal{S}_6 = \{ \hat{u}, \hat{v}, \hat{w}, 1 - \hat{u}, 1 - \hat{v}, 1 - \hat{w}, \hat{y}_a, \hat{y}_b, \hat{y}_w \}$
→ 1 for $\Delta = 0$
- $\rightarrow \mathcal{S}'_6 = \{ \hat{a} = \frac{\hat{u}}{\hat{v}\hat{w}}, \hat{b} = \frac{\hat{v}}{\hat{w}\hat{u}}, \hat{c} = \frac{\hat{w}}{\hat{u}\hat{v}}, \hat{d} = \frac{1-\hat{u}}{\hat{u}}, \hat{e} = \frac{1-\hat{v}}{\hat{v}}, \hat{f} = \frac{1-\hat{w}}{\hat{w}} \}$

- Kinematic map on letters:

$$\sqrt{\hat{a}} = d, \quad \hat{d} = a, \quad \text{plus cyclic relations}$$

$$S[A_6^{(1)}] = \left(-\frac{1}{2}\right) \hat{b} \otimes \hat{d} + \text{dihedral}$$


$$S[A_6^{(2)}] = \hat{b} \otimes \hat{d} \otimes \hat{d} \otimes \hat{d} + \frac{1}{2} \hat{b} \otimes \hat{b} \otimes \hat{b} \otimes \hat{d} + \text{dihedral}$$

...

L	number of terms
1	6
2	12
3	636
4	11,208
5	263,880
6	4,916,466
7	92,954,568
8	1,671,656,292

- Works through 7 loops!

Map covers entire phase space for 3-gluon form factor




- Soft is dual to collinear; collinear is dual to soft
- White regions in (u, v) map to some of $\hat{u}, \hat{v}, \hat{w} > 1$

Many special dual points

There is an “ f ” alphabet at all of these points, which is a way of writing multiple zeta values (MZV’s) so that the coaction is manifest.

F. Brown, 1102.1310;
O. Schnetz,
HyperlogProcedures



	$(\hat{u}, \hat{v}, \hat{w})$	(u, v, w)	functions
∇	$(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$	$(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$	$\sqrt[6]{1}$
\square	$(\frac{1}{2}, \frac{1}{2}, 0)$	$(0, 0, 1)$	$\text{Li}_2(\frac{1}{2}) + \text{logs}$
\bullet	$(1, 1, 1)$	$\lim_{u \rightarrow \infty} (u, u, 1-2u)$	MZVs
\circ	$(0, 0, 1)$	$(\frac{1}{2}, \frac{1}{2}, 0)$	MZVs + logs
\triangle	$(\frac{3}{4}, \frac{3}{4}, \frac{1}{4})$	$(-1, -1, 3)$	$\sqrt[6]{1}$
\boxplus	(∞, ∞, ∞)	$(1, 1, -1)$	alternating sums
\otimes	$\lim_{\hat{v} \rightarrow \infty} (1, \hat{v}, \hat{v})$	$\lim_{v \rightarrow \infty} (1, v, -v)$	MZVs
---	$(1, \hat{v}, \hat{v})$	$\lim_{v \rightarrow \infty} (u, v, 1-u-v)$	$\text{HPL}\{0, 1\}$
---	$(\hat{u}, \hat{u}, (1-2\hat{u})^2)$	$(u, u, 1-2u)$	$\text{HPL}\{-1, 0, 1\}$

The simplest point

- $(\hat{u}, \hat{v}, \hat{w}) = (1,1,1) \iff u, v \rightarrow \infty$

- At this point,

$$A_6^{(1)}(\cdot) = 0$$

$$F_3^{(1)}(\cdot) = 8\zeta_2$$

$$A_6^{(2)}(\cdot) = -9\zeta_4$$

$$F_3^{(2)}(\cdot) = 31\zeta_4$$

$$A_6^{(3)}(\cdot) = 121\zeta_6$$

$$F_3^{(3)}(\cdot) = -145\zeta_6$$

$$A_6^{(4)}(\cdot) = 120f_{3,5} - 48\zeta_2f_{3,3} - \frac{6381}{4}\zeta_8$$

$$F_3^{(4)}(\cdot) = 120f_{5,3} + \frac{11363}{4}\zeta_8$$

$$A_6^{(5)}(\cdot) = -2688f_{3,7} - 1560f_{5,5} + \mathcal{O}(\pi^2)$$


$$F_3^{(5)}(\cdot) = -2688f_{7,3} - 1560f_{5,5} + \mathcal{O}(\pi^2)$$

$$A_6^{(6)}(\cdot) = 48528f_{3,9} + 37296f_{5,7} + 21120f_{7,5} + \mathcal{O}(\pi^2)$$

$$F_3^{(6)}(\cdot) = 48528f_{9,3} + 37296f_{7,5} + 21120f_{5,7} + \mathcal{O}(\pi^2)$$

- Reversing ordering of words in f -alphabet, the blue values show that antipodal duality holds at these points beyond symbol level, modulo $i\pi$
- modulo $i\pi$ seems to be the best we can get from the antipode

Euclidean Region numerics





Bootstrap boundary data: Flux tubes at finite coupling

Alday, Gaiotto, Maldacena, Sever, Vieira, 1006.2788;


Basso, Sever, Vieira, 1303.1396, 1306.2058, 1402.3307, 1407.1736, 1508.03045

BSV+Caetano+Cordova, 1412.1132, 1508.02987



- Tile n -gon with pentagon transitions.
- Quantum integrability \rightarrow compute pentagons **exactly** in 't Hooft coupling
- 4d S-matrix as expansion (OPE) in **number of flux-tube excitations** = expansion around **near collinear limit**

The new FFOPE



- Form factors are Wilson loops in a **periodic** space, due to injection of operator momentum

Alday, Maldacena, 0710.1060; Maldacena, Zhiboedov, 1009.1139;
Brandhuber, Spence, Travaglini, Yang, 1011.1899

- Besides **pentagon transitions** \mathcal{P} , this program needs an **additional ingredient**, the **form factor transition** \mathcal{F}

Sever, Tumanov, Wilhelm, 2009.11297, 2105.13367, 2112.10569

OPE representation

- 6-gluon amplitude:

$$\mathcal{W}_{\text{hex}} = \sum_{\mathbf{a}} \int d\mathbf{u} P_{\mathbf{a}}(0|\mathbf{u}) P_{\mathbf{a}}(\bar{\mathbf{u}}|0) e^{-E(\mathbf{u})\tau + ip(\mathbf{u})\sigma + im\phi}$$

$$T = e^{-\tau}, S = e^{-\sigma}, F = e^{i\phi}. \quad v = \frac{T^2}{1+T^2} \rightarrow 0,$$

weak-coupling, $E = k + \mathcal{O}(g^2) \rightarrow$ expansion in T^k

- 3-gluon form factor: $\psi = \text{helicity 0 pairs of states}$

$$\mathcal{W}_3 = \sum_{\psi} e^{-E_{\psi}\tau + ip_{\psi}\sigma} \mathcal{P}(0|\psi) \mathcal{F}(\psi)$$

weak-coupling \rightarrow expansion in T^{2k} (no azimuthal angle ϕ)

OPE parametrizations

- Amplitude:
$$\hat{u} = \frac{1}{1 + (\hat{T} + \hat{S}\hat{F})(\hat{T} + \hat{S}/\hat{F})},$$

($\hat{F} = 1$ for $\Delta = 0$)

$$\hat{v} = \hat{u}\hat{w}\hat{S}^2/\hat{T}^2, \quad \hat{w} = \frac{\hat{T}^2}{1 + \hat{T}^2}$$

- Form factor:
$$u = \frac{1}{1 + S^2 + T^2}, \quad v = \frac{T^2}{1 + T^2},$$

$$w = \frac{1}{(1 + T^2)(1 + S^{-2}(1 + T^2))},$$

- Apply kinematic map \rightarrow
$$\hat{T} = \frac{T}{S}, \quad \hat{S} = \frac{1}{TS}$$

- Apparently some correspondence between **single** flux tube excitations for the amplitude (T^1) and **double** (or bound state) excitations for the form factor (T^2)

8-gluon Amp \leftrightarrow 4-gluon FF

LD, Ö. Gürdoğan, Y.-T. Liu A. McLeod, M. Wilhelm, in progress

- We have a **candidate kinematic map** for a **4-dimensional** surface (4-gluon FF is 5d).
- $\mathcal{S}[R_8^{(2)}]$ is known [S. Caron-Huot, 1105.5606](#)
- The **kinematic+antipodal** maps take it to a symbol with 40 letters, the first 8 of which are “right”:
$$u_i = \frac{s_{i,i+1}}{s_{1234}}, \quad v_i = \frac{s_{i,i+1,i+2}}{s_{1234}}$$
- But we still have to run more checks on this **candidate 2-loop 4-gluon form factor**

8-4 Kinematic Map in OPE Parametrization

- 8-point amplitude has D_8 dihedral symmetry; change it to D_4 of the form factor by requiring



$$\hat{T}_3 = \hat{T}_1, \quad \hat{S}_3 = \hat{S}_1, \quad \hat{F}_3 = \hat{F}_1$$

- To get $\mathcal{S}[R_8^{(2)}]$ to have only 8 final entries, we also fix $\hat{F}_1 = \hat{F}_2 = 1$.

- The kinematic map becomes

$$\hat{T}_1 = \frac{T}{S}, \quad \hat{S}_1 = \frac{1}{TS},$$

$$\hat{T}_2 = \frac{T_2}{S_2}, \quad \hat{S}_2 = \frac{1}{T_2 S_2} \quad \text{and requires } F_2 = i$$



Summary & Outlook

- Form factors as well as scattering amplitudes in planar $N=4$ SYM can now be **bootstrapped** to high loop order
- Remarkably simple behavior at “origin”
- Comparing the 3-gluon form factor to the 6-gluon amplitude, a strange new antipodal duality swaps the role of branch cuts and derivatives, and may map single flux-tube excitations (amplitude) to doubles (form factor).
- What is the underlying physical reason for this duality?
- (How) does it hold at strong coupling?
- (How much) can we verify of it at the 8-4 level, and beyond?
- How much can we exploit it to learn more about both amplitudes and form factors?

Extra Slides


Removing Amplitude (or Form Factor) Infrared Divergences

- On-shell amplitudes **IR divergent** due to long-range gluons

- Polygonal Wilson loops **UV divergent** at cusps,
anomalous dimension Γ_{cusp}

– known to all orders in planar N=4 SYM:

Beisert, Eden, Staudacher, hep-th/0610251



- Both removed by dividing by **BDS-like ansatz**

Bern, LD, Smirnov, hep-th/0505205, Alday, Gaiotto, Maldacena, 0911.4708

- Normalized [MHV] amplitude is finite, dual conformal invariant, also **uniquely** (up to **constant**) maintains important symbol adjacency relations due to causality (Steinmann relations for **3-particle invariants**):

$$\mathcal{E}(u_i) = \lim_{\epsilon \rightarrow 0} \frac{\mathcal{A}_6(s_{i,i+1}, \epsilon)}{\mathcal{A}_6^{\text{BDS-like}}(s_{i,i+1}, \epsilon)} = \exp\left[\frac{\Gamma_{\text{cusp}}}{4} \mathcal{E}^{(1)} + R_6\right]$$

↑
remainder function

BDS & BDS-like normalization for \mathcal{F}_3

$$\frac{\mathcal{F}_3}{\mathcal{F}_3^{\text{MHV, tree}}} = \exp \left\{ \sum_{L=1}^{\infty} g^{2L} \left[\left(\frac{\Gamma_{\text{cusp}}^{(L)}}{4} + \mathcal{O}(\epsilon) \right) M^{1\text{-loop}}(L\epsilon) + C^{(L)} + R^{(L)}(u, v, w) \right] \right\}$$

BDS ansatz

remainder function only a function of u, v, w ; vanishes in all collinear limits, but no adjacency constraints

split 1-loop amplitude judiciously:

$$\frac{\mathcal{F}_3^{1\text{-loop}}}{\mathcal{F}_3^{\text{MHV, tree}}} \equiv M^{1\text{-loop}}(\epsilon) = M(\epsilon) + \mathcal{E}^{(1)}(u, v, w)$$

$$M(\epsilon) = -\frac{1}{\epsilon^2} \sum_{i=1}^3 \left(\frac{\mu^2}{-s_{i,i+1}} \right)^\epsilon - \frac{7}{2} \zeta_2 + \frac{3}{\epsilon}$$

\mathcal{E} obeys "adjacency constraints"

$$\mathcal{E}^{(1)}(u, v, w) = \left[\text{Li}_2\left(1 - \frac{v}{w}\right) + \text{Li}_2\left(1 - \frac{1}{w}\right) \right] \quad \mathcal{E}^{(1),u} + \mathcal{E}^{(1),1-u} = 0$$

Now divide by $\mathcal{F}_3^{\text{MHV, tree}}$


$$\frac{\mathcal{F}_3^{\text{BDS-like}}}{\mathcal{F}_3^{\text{MHV, tree}}} = \exp \left\{ \sum_{L=1}^{\infty} g^{2L} \left[\left(\frac{\Gamma_{\text{cusp}}}{4} + \mathcal{O}(\epsilon) \right) M(L\epsilon) + C^{(L)} \right] \right\} \Rightarrow \mathcal{E} = \exp \left[\frac{\Gamma_{\text{cusp}}}{4} \mathcal{E}^{(1)} + R \right]$$

Number of (symbol-level) linearly independent $\{n, 1, \dots, 1\}$ coproducts ($2L - n$ derivatives)

weight n	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$L = 1$	1	3	1														
$L = 2$	1	3	6	3	1												
$L = 3$	1	3	9	12	6	3	1										
$L = 4$	1	3	9	21	24	12	6	3	1								
$L = 5$	1	3	9	21	46	45	24	12	6	3	1						
$L = 6$	1	3	9	21	48	99	85	45	24	12	6	3	1				
$L = 7$	1	3	9	21	48	108	236	155	85	45	24	12	6	3	1		
$L = 8$	1	3	9	21	48	108	242	466	279	155	85	45	24	12	6	3	1

- Properly normalized L loop N=4 form factors $\mathcal{E}^{(L)}$ belong to a small space \mathcal{C} , dimension saturates on left
- $\mathcal{E}^{(L)}$ also obeys multiple-final-entry relations, saturation on right

Some numerics





I = decay / Euclidean

IIa,b,c = scattering / spacelike operator


IIIa,b,c = scattering / timelike operator

Real “impact factor” appears in space-like Regge limit, $\nu \rightarrow \infty$

Remainder function R is nontrivial
function of $u = \frac{s_{12}}{m_H^2}$ as $s_{23} \rightarrow \infty$



Numerical implications of antipodal duality?



Values of HPLs $\{0,1\}$ at $u = 1$

- Classical polylogs

$$\text{Li}_n(u) = \int_0^u \frac{dt}{t} \text{Li}_{n-1}(t) = \sum_{k=1}^{\infty} \frac{u^k}{k^n}$$

evaluate to Riemann zeta values

$$\text{Li}_n(1) = \sum_{k=1}^{\infty} \frac{1}{k^n} = \zeta(n) \equiv \zeta_n$$

- HPL's evaluate to **nested sums** called **multiple zeta values (MZVs)**:

$$\zeta_{n_1, n_2, \dots, n_m} = \sum_{k_1 > k_2 > \dots > k_m > 0} \frac{1}{k_1^{n_1} k_2^{n_2} \dots k_m^{n_m}}$$

Weight $n = n_1 + n_2 + \dots + n_m$

- **MZV's** obey many identities, e.g. stuffle

$$\zeta_{n_1} \zeta_{n_2} = \zeta_{n_1, n_2} + \zeta_{n_2, n_1} + \zeta_{n_1 + n_2}$$

- All reducible to Riemann zeta values until **weight 8**.

Irreducible MZVs: $\zeta_{5,3}, \zeta_{7,3}, \zeta_{5,3,3}, \zeta_{9,3}, \zeta_{6,4,1,1}, \dots$

Many “empirical” adjacency constraints


$$F^{d,e} = F^{e,d} = F^{e,f} = F^{f,e} = F^{f,d} = F^{d,f} = 0$$

Hold for 2 loop QCD amplitudes too, planar and nonplanar!

LD, Mcleod, Wilhelm, 2012.12286

$$F^{a,d} = F^{d,a} = F^{b,e} = F^{e,b} = F^{c,f} = F^{f,c} = 0$$

Latter are NEW: Hold for planar N=4 SYM to 8 loops!



Mnemonic for dihedral symmetry;
6 dashed lines indicate 12 forbidden pairs.

Empirical multi-final entry relations

1. $\xi^a = 0$ (plus dihedral images)

2. $\xi^{a,e} = \xi^{a,f}$ (plus ...)

3. $\xi^{a,b,d} = 0, \quad \xi^{a,e,e} = -\xi^{a,f,f},$
 $\xi^{e,a,f} = \xi^{f,a,f} - \xi^{a,f,f}$

4.

Number of remaining parameters in form-factor ansatz after imposing constraints

L	2	3	4	5	6	7	8
symbols in \mathcal{C}	48	249	1290	6654	34219	????	????
dihedral symmetry	11	51	247	1219	????	????	????
$(L - 1)$ final entries	5	9	20	44	86	???	???
L^{th} discontinuity	2	5	17	38	75	???	??
collinear limit	0	1	2	8	19	70	6
OPE $T^2 \ln^{L-1} T$	0	0	0	4	12	56	0
OPE $T^2 \ln^{L-2} T$	0	0	0	0	0	36	0
OPE $T^2 \ln^{L-3} T$	0	0	0	0	0	0	0
OPE $T^2 \ln^{L-4} T$	0	0	0	0	0	0	0
OPE $T^2 \ln^{L-5} T$	0	0	0	0	0	0	0

Table 4: Number of parameters left when bootstrapping the form factor $\mathcal{E}^{(L)}$ at L -loop order in the function space \mathcal{C} at symbol level, using all the conditions on the final $(L - 1)$ entries, which can be deduced at $(L - 1)$ loops.

The [Dual] Conformal Group


$SO(4,2) \supset SO(3,1)$ [rotations+boosts] + translations+dilatations + special-conformal

$$15 = 3 + 3 + 4 + 1 + 4$$

- The nontrivial generators are special conformal K^μ
- Correspond to inversion · translation · inversion
- To obtain a [dual] conformally invariant function $f(x_{ij}^2)$ just have to check invariance under inversion,

$$x_i^\mu \rightarrow x_i^\mu / x_i^2$$

Different routes to perturbative amplitudes



Beyond 8-4

- The map $\hat{T}_1 = \frac{T}{S}$, $\hat{S}_1 = \frac{1}{TS}$, $\hat{T}_2 = \frac{T_2}{S_2}$, $\hat{S}_2 = \frac{1}{T_2 S_2}$

seems **likely** to generalize to give rise to a $2(n - 2)$ parameter subspace of the full $3n - 7$ dimensional n -point form factor kinematics, presumably from setting $F_2 = \dots = F_{n-2} = i$

- We can **conjecture** that **antipodal duality** applies on this subspace
- But there is still a lot to be checked!